

AD \_\_\_\_\_

GRANT NO: DAMD17-94-J-4021


TITLE: Spatial Distribution of the EGF Receptor System in the Regulation of Breast Epithelia Cell Growth and Organization

PRINCIPAL INVESTIGATOR(S): Patrick Burke

CONTRACTING ORGANIZATION: University of Utah  
Graduate School  
Salt Lake City, Utah 84112

REPORT DATE: June 12, 1995

TYPE OF REPORT: Annual



PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19950901 041

DTIC QUALITY INSPECTED 6

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 12, 1995	3. REPORT TYPE AND DATES COVERED Annual May 1994-May 1995	
4. TITLE AND SUBTITLE Spatial Distribution of the EGF Receptor System in the Regulation of Breast Epithelia Cell Growth and Organization			5. FUNDING NUMBERS  DAMD17-94-J-4021	
6. AUTHOR(S)  Patrick Burke				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Utah Graduate School Salt Lake City, Utah 84112			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  184A1 human mammary epithelial cells (HMECs) form organotypic structures when plated on Matrigel. However, the organotypic structures differentiate into squamous, not glandular, epithelia. They can not be used to examine the functional significance of a polarized epidermal growth factor receptor system in a glandular epithelia. This experimental system may be useful for investigating cell-extracellular matrix interactions that control EGF-R expression. 184A1s and subclones of the 184A1 cell line express proteins necessary for structural and functional polarization on Transwells, but do not assemble these proteins into tight junctions. MCF-7 cells express ZO-1 and assemble the protein into tight junctions when plated onto Transwells. MCF-7s also segregate B-1 integrin into the basal-lateral membrane compartment. MCF-7 cells respond mitogenically to EGF-R occupancy and may provide an experimental system with which to address the primary aim of my proposal: to determine if the loss of the correct spatial organization or inappropriate expression of the EGF-R system provides a growth advantage, enhances cell motility, or ochanges the differentiated state of normally organized epithelial cells.				
14. SUBJECT TERMS EGF-Receptor; Breast Cancer; Mammary Epithelial Cells; Polarized Epithelia; Differentiation; Matrigel			15. NUMBER OF PAGES 10	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

X Where copyrighted material is quoted, permission has been obtained to use such material.

X Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

X Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

X For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

X In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

X In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

X In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
HTIS GRAAI	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

*Robert J. Burke* 6/12/95  
PI - Signature Date

## Table of Contents

### Page Number(s):

1	Front Cover
2	Form SF298
3	Foreword
4	Table of Contents
5	Introduction
6	Body
7	Conclusions
8	References
9-10	Appendices: Figures 1-3

## INTRODUCTION:

### Spatial Distribution of the EGF Receptor System in the Regulation of Breast Epithelial Cell Growth and Organization

The epidermal growth factor receptor (EGF-R) system is necessary for the motility, proliferation and differentiation of human mammary epithelial cells (HMECs) *in vitro*. Additionally, the EGF-R system displays a highly organized spatial distribution *in vivo*. Because the EGF-R system plays a central role in HMEC proliferation, it is reasonable to suspect that any defects in its regulation could lead to the clonal expansion of 'pre-malignant' cell populations. Such expanding clonal populations could give rise to cancerous clones. The spatial distribution of the EGF-R system is highly organized *in vivo*. *In vivo*, the receptor and one of its ligands, transforming growth factor alpha (TGF-  $\alpha$ ), are localized to the basolateral surface of mammary alveolar structures; on the other hand, epidermal growth factor (EGF) is synthesized and secreted from the apical side into the lumen of the alveoli. Regulation of the EGF-R system could be disrupted by removing the spatial restrictions which segregate one of the ligands from the receptor and/or the receptor from second messenger systems. I will investigate the spatial distribution of both the EGF-R and its ligands during the organization of HMECs on basement membranes (BM) *in vitro* and determine the consequences of disorganizing this distribution.

#### SPECIFIC AIMS:

1. Define the spatial distribution and expression levels of the EGF-R system in proliferating and spatially organized normal HMECs.
2. Determine whether a loss of the correct spatial organization or inappropriate expression of the EGF-R system provides a growth advantage, enhances cell motility, or changes the differentiated state of normally organized epithelial cells.

## BODY OF REPORT

The HMECs I chose to use in this work are designated 184A1 and were isolated by Martha Stampfer as an immortalized derivative from the parental 184 cell line (1). They exhibit the ability to organize into organotypic structures when plated onto the extracellular matrix material Matrigel (extracted from Englebreth-Holm-Swarm murine sarcoma), previously shown in proposal. Histological sections of the organotypic structures stained with hematoxylin and eosin strongly suggest that after 3 weeks in culture on matrigel the 184A1s differentiate into squamous, rather than glandular epithelia as initial sections suggested. Identification of intracellular bridging, layering of cells and the secretion of cytokeratins are the ultrastructural basis upon which the organotypic structures have been identified as squamous in nature.

EGF-R levels have been examined during this squamous differentiation process and appear to decrease rapidly upon contact with the basement membrane and continue to decrease over a seven day period (figure 1). The mechanism of this regulation is unknown, but will be investigated further to determine if it is post-translational, post transcriptional or both. The involvement of specific integrin-extracellular matrix interactions in this process will also be investigated.

184A1s and subclones of 184A1s were plated onto tissue culture membranes as a way to induce epithelial polarization. Using membranes is a common experimental approach by which to investigate properties of polarized epithelia. 184A1s express the protein ZO-1, an integral component of the tight junctions that is characteristic of polarized epithelia; however when compared to the positive control, polarized MDCK cells, it is obvious that the protein is not part of a well formed intercellular junction (figure 2) (2,3,4). Different media and substratum conditions were used to induce tight junction formation, but no attempts have been successful thus far.

I have investigated the ability of a non-aggressive breast cancer cell line MCF-7 to functionally polarize as an alternative model to address my initial specific aims. The human colon cancer cell line, Caco-2, has served as an analogous system for those studying the structure and function of polarized intestinal epithelia (3,4). Initial immunofluorescence results strongly suggest that MCF-7 cells polarize when plated onto tissue culture membranes (Costar, Transwells) as determined by the distribution of ZO-1 and Beta-1 integrin. The distribution of both is consistent with a polarized epithelia (figure 3). MCF-7 cells are a classic example of an EGF-R autocrine cell; their proliferation is estrogen dependent which imparts much of its growth influence through induction of the EGF-R/TGF- $\alpha$  autocrine loop (5,6,7). MCF-7 cells may serve as an effective model for studying the functional importance of a polarized distribution of the EGF-R system.

## CONCLUSIONS

1. Organotypic HMECs structures differentiate into squamous, not glandular, epithelia when plated on matrigel. This experimental approach may be useful for investigating cell-extracellular matrix interactions that control EGF-R expression. The mechanism will be determined to be post-translational and/or transcriptional, and further investigations will examine the role of laminin and  $\beta$ -1 integrin in this process.
2. 184A1s and subclones express proteins necessary for structural and functional polarization on transwells, but do not assemble these proteins into tight junctions.
3. MCF-7 cells express ZO-1 and assemble the protein into tight junctions when plated onto transwells. MCF-7s also segregate  $\beta$ -1 integrin into the basal-lateral membrane compartment. Further studies, including electron microscopy, immunofluorescence and measurements of electrical resistance will better define the degree of structural and functional polarization in MCF-7s.


## References

1. Stampher, M.R. and Bartley, J.C. (1985) Induction of Transformation and Continuous Cell Lines from Normal Human Mammary Epithelial Cells After Exposure to Benzo(a)pyrene. *Proc. Natl. Acad. Sci. U.S.A.* **82**: 2394-2401.
2. Anderson, J.M., Balda, M.S., Fanning, A.S. (1993) The Structure and Regulation of Tight Junctions. *Curr. Opin. Cell Biol.* **5**: 772-778.4.
3. Nelson, W.J. (1992) Regulation of Cell Surface Polarity from Bacteria to Mammals. *Science*. **258**: 948-954.
4. Rodriguez-Boulán, E. and Nelson, W.J. (1989) Morphogenesis of the Polarized Epithelial Cell Phenotype. *Science*. **245**: 718-725.
5. Colomb, E., Berthon, P., Dussert, C., Calvo, F. and Martin, P.M. (1991) Estradiol and EGF requirements for cell-cycle progression of normal human mammary epithelial cells in culture. *Int J Cancer*. **49**: 932-7.
6. Kern, F.G., Chevelle, A.L. and Liu, Y. (1990) Growth Factor Receptors and the Progression of Breast Cancer. *Semin. Cancer Biol.* **1**: 317-328.
7. Bates, S.E., Davidson, N.E., Valverius, E.M., Freter, C.E., Dickson, R.B., Tam, J.P., Kudlow, J.E., Lippman, M.E., Salomon, D.S. (1988) Expression of Transforming Growth Factor  $\alpha$  and its Messenger Ribonucleic Acid in Human Breast Cancer: Its Regulation by Estrogen and its Possible Functional Significance. *Mol. Endocrin.* **2**: 543-555.




## Appendix

**Figure 1:** Decreased EGF-R Expression During Organization on Matrigel as determined by western blot, equal amounts of protein were loaded in each lane; the control is cells grown on plastic



**Figure 2:** 184 HMECs Do Not Form Well Differentiated Polarized Epithelial Layers on Membranes



## Appendix

**Figure 3:** Polarization of MCF-7, Compare to figure 2 and the localization of ZO-1 and EGF-R in MDCKs (positive controls)

